



Open Access  
Scan to access more  
free content

# Controlled rehabilitative and supportive care intervention trials in patients with high-grade gliomas and their caregivers: a systematic review

K Piil,<sup>1,2</sup> M Juhler,<sup>1,3</sup> J Jakobsen,<sup>4</sup> M Jarden<sup>2</sup>

## ABSTRACT

**Background** Patients diagnosed with high-grade gliomas experience a varying and complex symptom burden, and face a high mortality rate. As a consequence, patients with high-grade gliomas and their caregivers have imminent and changing rehabilitative and supportive care needs.

**Objectives** To give a detailed overview of non-pharmacological rehabilitative and supportive care interventions for patients with high-grade gliomas and/or their caregivers, and provide an appraisal of the methodological quality of these studies.

**Method** PubMed, Cumulative Index of Nursing and Allied Health Literature and Embase were searched for literature published from 1995 to May 2013. Data from eight studies were reviewed for substantive methods and results. Methodological quality was described and assessed using the scoring system for appraising mixed methods research and concomitantly appraising qualitative, quantitative and mixed methods primary studies in mixed study reviews.

**Results** The search yielded 914 unique publications, of which 9 were classified eligible for this review. There is preliminary evidence that cognitive group therapy improves memory skills in patients with high-grade gliomas, early physical training improves functional outcome and massage therapy reduces stress. Patients and caregivers found that telephone follow-up and a specialist nurse function was an effective and useful way to achieve information and support. Finally, psycho-education increased feelings of mastery among caregivers.

**Conclusions** As evidence is beginning to emerge, there is a need for well-designed longitudinal and randomised controlled trials of non-pharmacological interventions in high-grade

glioma patients and their caregivers in order to develop clinical guidelines for supportive and rehabilitative approaches in this unique population.

## INTRODUCTION

High-grade glioma (HGG) is the most malignant type of brain tumour (BT) occurring most frequently in people aged 45–70.<sup>1</sup> Treated with optimal therapy, the median survival is 12–15 months with a 5-year survival of 10%.<sup>2</sup> Patients with HGG experience a high symptom burden related to the disease and the surgical procedures and medical treatments.<sup>3–4</sup> Different types of cerebral symptoms and complications may occur, including global cerebral symptoms (fatigue, nausea, headache, confusion) focal symptoms (hemiparesis, seizures, speech difficulties), neurocognitive deficits (impaired attention, concentration difficulties, reduced short-term memory, personality changes) and emotional symptoms (depression, anxiety, stress).<sup>5–9</sup> As a consequence, patients with HGG and their caregivers have a number of rehabilitative and supportive care needs. The literature recommends that future intervention studies include individually tailored communication and specialist support<sup>10–12</sup> to relieve emotional distress in patients with gliomas<sup>13</sup> and stress among caregivers.<sup>14</sup> However, little attention has been paid to non-pharmacological rehabilitative and supportive care interventions targeted patients with HGG and/or their caregivers.<sup>15</sup> A recent Cochrane literature review<sup>16</sup> was unable to identify any randomised controlled trials (RCT) or controlled clinical trials of multidisciplinary

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bmjspcare-2013-000593>).

<sup>1</sup>Department of Neurosurgery, The University Hospital of Copenhagen, Rigshospitalet, Copenhagen, Denmark

<sup>2</sup>The University Hospitals Center for Health Research (UCSF), Center for Integrated Rehabilitation for Patients with Cancer (CIRE), Copenhagen, Denmark

<sup>3</sup>Department of Clinical Medicine, Section of Neurology, Psychiatry and Sensory Sciences, The University of Copenhagen, Copenhagen K, Denmark

<sup>4</sup>The University Hospital of Copenhagen; Rigshospitalet, Neuroscience Center, Copenhagen, Denmark

## Correspondence to

Dr Karin Piil, Department of Neurosurgery, University Hospital of Copenhagen; Rigshospitalet, UCSF 9701, Blegdamsvej 9, Copenhagen 2100, Denmark; [Karin.piil@rh.regionh.dk](mailto:Karin.piil@rh.regionh.dk)

Received 12 September 2013

Revised 11 April 2014

Accepted 30 April 2014

Published Online First

2 June 2014



CrossMark

**To cite:** Piil K, Juhler M, Jakobsen J, et al. *BMJ Supportive & Palliative Care* 2016;**6**:27–34.

rehabilitation interventions in adults after primary BT treatment.<sup>16</sup> The majority of studies include participants with different subtypes of BT; however, tumour progression varies depending on tumour type. HGG is a rapidly progressive life-threatening disease, while other subgroups of BT progress more slowly. This is supported by investigators who underpin the importance of research on potential differences in needs among the heterogeneous population of patients with BT.<sup>17 18</sup> Presently, there are no evidence-based rehabilitative and supportive care guidelines that aim to reduce loss of cognition, function and quality of life (QOL). To our knowledge, this is the first systematic literature review that gives a detailed overview of rehabilitative and supportive care intervention studies in patients with HGG and/or their caregivers with a systematic appraisal of the methodological quality of each publication.

## METHODS

This systematic literature search reviews non-pharmacological rehabilitative and supportive care interventions in patients with HGG and their caregivers. Data regarding population, study design, intervention and main results are extracted, synthesised and discussed. The selected publications are appraised for methodological quality. Lastly, implications for future research and conclusions are presented.

### Literature search

The study population includes patients with HGG and/or their caregivers in non-pharmacological rehabilitative and supportive care intervention studies. Searches were performed in PubMed, Cumulative Index of Nursing and Allied Health Literature (CINAHL) and Embase. Combinations of the following 'Medical Subject Heading' (MeSH) terms and free text terms were used: 'brain cancer', 'brain neoplasms', 'brain tumor', 'high-grade glioma', 'malignant glioma', 'rehabilitation', 'training', 'expectations', 'needs', 'quality of life', 'supportive care', 'intervention', 'program', 'intervention studies', 'caregiver', 'spouse', 'relative' and 'next of kin'. The search strategy is illustrated in figure 1. The full search strategy including search terms used for each database is available and may be requested by contacting the corresponding author (KP). Reference Manager Software program was used to identify duplicates. The search was limited to English-, Danish-, Norwegian- and Swedish-language publications published from 1995 to May 2013. The systematic literature search was guided by the PRISMA guidelines—preferred reporting items for systematic reviews and meta-analyses developed by Cochrane Collaboration.<sup>19</sup>

### Inclusion and exclusion criteria

Publications included are randomised and non-randomised controlled trials of unidisciplinary or multidisciplinary non-pharmacological rehabilitative and/or supportive care interventions in patients with

HGG and/or their caregivers. Studies with different subtypes of BT are included only if statistical analysis was performed in the HGG population. Trials studying medical and pharmaceutical treatments are excluded from this review.

### Data extraction and quality assessment

Information regarding the population (sample size, age and attrition), the study design (focus and time of enrolment), the intervention (context, type, mode, duration, frequency, length and adherence) and main results were extracted and systemised using NVivo software program (see online supplementary table S1).<sup>20</sup> Assessment of methodological quality and scoring of each publication was carried out independently by two researchers (KP and MJA). There was no discrepancy between the raters' scores. The online supplementary table S2 shows an evaluation of the methodological quality of each publication. Methodological quality was assessed using the scoring system for appraising mixed methods research and concomitantly appraising qualitative, quantitative and mixed methods primary studies in mixed study reviews.<sup>21</sup> Methodological quality for the qualitative studies was assessed for the presence or absence of the following criteria: a qualitative objective or question, appropriate qualitative approach or design or method, a description of the context, a description of participants and justification of sampling, a description of qualitative data and a discussion of researchers' reflexivity collection and analysis. Methodological quality for the quantitative experimental studies was appraised for the presence or absence of the following criteria: an appropriate sequence generation and/or randomisation, allocation concealment and/or blinding, and presentation of complete outcome data and/or low withdrawal/dropout. For each criterion, the presence or absence is scored 1 or 0, respectively. The quality score is calculated as a percentage: (number of 'presence' responses divided by the number of relevant criteria)×100. A further descriptive and detailed critical appraisal of methodological quality is presented.

### Ethics

This systematic review is part of a larger study, which is registered at the Research Ethics Committees at the Regional capital in Copenhagen, Denmark (33460), and the Danish Data Protection Agency (2007-58-0015/30-0758).

## RESULTS

A systematic search was carried out by an information specialist and the primary investigator (KP) in June 2013. The search resulted in 748 hits in PubMed, 114 hits in CINAHL and 52 hits in Embase and is schematically outlined in figure 1. No duplicates were identified, resulting in 914 unique titles. The titles




Figure 1 PRISMA flow diagram.

and/or abstracts of the 914 publications were reviewed by the primary investigator (KP), and 894 publications were excluded. Two researchers (KP and MJA) reviewed 20 full manuscripts, which appeared potentially relevant, and 11 were excluded. Nine publications describing eight different interventions met the inclusion criteria.<sup>22–30</sup> Sardell *et al*'s<sup>26</sup> mixed BT population study was included because all 45 patients (n=43 HGG; n=1 primitive neuroectodermal tumour; n=1 oligoastrocytoma) received primary therapy for high-grade malignancy.

### Study characteristics

The online supplementary table S1 describes populations studied, interventions employed and main results.

The nine publications were published in seven different scientific journals between 2000 and 2012, the majority being published within the last 5 years. The time of patient and/or caregiver enrolment to the interventions varied in the studies, from time of

diagnosis to late stages of the disease. Four studies were initiated at time of diagnosis,<sup>24 25 27 28 30</sup> one after hospital discharge<sup>23</sup> and two studies 15–16 months after diagnosis.<sup>22 26</sup> All interventions were supervised, seven were carried out individually<sup>23–30</sup> and one intervention was a group intervention.<sup>22</sup> Six studies were unidisciplinary<sup>22 23 26–30</sup> and two were multidisciplinary as they combined physical training with occupational and speech therapy, respectively.<sup>24 25</sup>

### Populations studied

The studies primarily included patients diagnosed with HGG. However, one publication included only caregivers,<sup>29</sup> and one included both patients and their caregivers.<sup>27 28</sup> This was the only intervention that continued to intervene with caregivers after the patient died.<sup>27 28</sup> The age of the patients ranged from 23 to 79 years (mean 55.8). The age of the caregivers was not reported. Sample sizes ranged from 11 to 121 participants (mean 50.8). The demographic characteristics of the participants were included in all but one

study<sup>30</sup> and not outlined according to subgroup diagnoses in two studies.<sup>23 24</sup>

### Intervention and outcome characteristics

Nine publications (eight interventions) reported the results of non-pharmacological rehabilitative and/or supportive care interventions in patients with HGG and/or their caregivers<sup>22–30</sup> (see online supplementary table S1). Hassler *et al*<sup>22</sup> examined the effect of supervised cognitive group training in patients with HGG, Pace *et al*,<sup>23</sup> Geler-Kulcu *et al*<sup>24</sup> and Bartolo *et al*<sup>25</sup> studied the effect of supervised physical training in patients with HGG and Sardell *et al*,<sup>26</sup> Spetz *et al*<sup>27 28</sup> and Boele *et al*<sup>29</sup> (four publications) tested nurse-led telephone follow-up (NTF), a specialist nurse (SN) function in patients and caregivers and psycho-education in caregivers, respectively. Keir<sup>30</sup> examined the effect of massage therapy among HGG patients.

Five interventions were conducted in a hospital setting,<sup>22 24 25 29 30</sup> one of the physical training programmes was carried out as a home programme<sup>23</sup> and two follow-up studies took place in a combination of a home and hospital setting.<sup>26–28</sup> The length of the interventions varied considerably, from 2 weeks to comprise the entire disease and treatment trajectory until post bereavement. The duration of the interventions was between 6 and 57 h (median 27.7);<sup>22 24 29 30</sup> however, it was not specified for three supportive care interventions.<sup>26–28</sup>

Hassler *et al*'s<sup>22</sup> cognitive group training programme found significant improvement in memory skills measured by verbal memory total learning ( $p=0.04$ ) and no change in verbal memory delayed recall ( $p=0.11$ ), psychomotor speed ( $p=0.22$ ), sustained attention ( $p=0.17$ ) and verbal fluency ( $p=0.29$ ). Pace *et al*<sup>23</sup> introduced a home rehabilitation programme with individual physical training and reported clinical improvements for the BT group on KPS and The Bartel Index (BI) measuring Activities of Daily Living. Among these patients, a significant improvement in median BI ( $p=0.001$ ) and median KPS ( $p=0.001$ ) was reported. Subgroup data showed no difference between histological subgroups; however, a functional gain (not specified) was observed in 26% of the HGG population ( $N=62$ ). Geler-Kulcu *et al*'s<sup>24</sup> trial with physical and occupational therapy compared BT patients postoperatively and patients with stroke. Both groups improved in Motor Assessment Scale (MAS) ( $p=0.025/p=0.002$ ), Postural Assessment Scale for Stroke (PASS) ( $p=0.002/p=0.002$ ), Berg Balance Scale (BBS) ( $p=0.011/p=0.109$ ) and Functional Impairment Measurement (FIM-mobility) ( $p=0.043/p=0.007$ ). No statistical difference between BT subgroups or between the group of BT patients and stroke patients was observed in any measurement (MAS  $p=0.718$ , PASS  $p=0.817$ , BBS  $p=0.109$ , FIM-mobility  $p=0.660$ ). Bartolo *et al*<sup>25</sup> found similar results in a

physical training and speech therapy intervention in a BT group and a stroke group. All outcome measures improved at postintervention for both groups ( $p<0.001$ ), though without statistical differences between groups. A subgroup analysis (meningioma, HGG and stroke) revealed that patients with meningioma achieved better independence in activities of daily living ( $p=0.02$ ) and mobility ( $p=0.04$ ) compared with the HGG and stroke group. However, the total FIM scores and other clinical scales revealed no differences between the subgroups. There were four different types of supportive care interventions.<sup>26–29</sup> Sardell *et al*<sup>26</sup> showed that a monthly and ongoing NTF of patients with HGG was an effective alternative to conventional clinic follow-up with face-to-face consultation. Patient satisfaction was high and the numbers of unscheduled clinic visits were minimised to include acute problems as the NTF assisted the patients with a variety of other problems by telephone. A qualitative study carried out by Spetz *et al*<sup>27 28</sup> included both patients and caregivers and showed that an SN role was a useful way to provide information and support. The SN function increased the family's autonomy and facilitated decision making within the family. The SN was a resource for the patients, the caregivers and the entire family. Boele *et al*'s<sup>29</sup> RCT found that a psycho-education intervention in caregivers during a 6-week period increased feelings of mastery ( $p=0.021$ ). Keir's 4-week massage therapy intervention reduced stress between weeks 2 and 3 ( $p=0.010$ ) and improved emotional well-being ( $p\leq 0.001$ ), BT concerns ( $p\leq 0.013$ ) and social/family well-being ( $p\leq 0.024$ ) at post-testing. All participants were below their baseline stress score 1 week after the massage intervention was completed.<sup>30</sup>

### Description of study methodological quality

Systematic methodological ratings of all the included nine papers, with respect to each relevant criterion, are shown in the online supplementary table S2. Spetz *et al*'s<sup>27 28</sup> intervention study applied a qualitative approach to explore the effect of the SN function. It fulfilled the methodological quality criteria by 100%. The study formulated a qualitative objective, applied an appropriate qualitative method, described the context, the participants and made a justification of sampling. Further, appropriate qualitative data collection and analysis was carried out and the researcher's reflexivity was described. Seven intervention studies applied quantitative experimental designs<sup>22–26 29 30</sup>; the neurocognitive training study,<sup>22</sup> the home rehabilitation programme,<sup>23</sup> the two physical training trials comparing BT patients and stroke patients,<sup>24 25</sup> the evaluation of NTF,<sup>26</sup> the psycho-education study<sup>29</sup> and the massage study.<sup>30</sup> All fulfilled 33.3% of the quality criteria as they only presented a complete outcome data and/or low dropout rate, except for the psycho-education study that scored 33.3%, as it



fulfilled the criteria of applying an appropriate randomisation.<sup>29</sup>

Further details regarding the study design, inclusion and exclusion criteria, risk of bias, attrition rates and adverse events are described in the following. All studies were designed as controlled interventions trials. One study published in two papers applied a qualitative design (interview, observation).<sup>27–28</sup> The remaining seven studies used quantitative designs (RCT, case-control, clinical controlled pilot trials, comparative trials).<sup>22–26–30</sup> All interventions carried out pre-post measurements.

Three interventions were conducted during the early stage of the disease.<sup>23–25</sup>

Statistical methods and p values were presented in all but two studies.<sup>26–28</sup> Validated measurement outcomes were applied in all studies, except the qualitative study.<sup>27–28</sup> All interventions are described in detail in such a way that would allow duplication of the intervention design, except the home training study.<sup>23</sup>

All nine papers defined an inclusion and exclusion criteria. Exclusions were often due to patients/caregivers speaking a language unfamiliar to the investigators or participants being unable to complete the questionnaires. Aside from these systematic exclusions, there was a range of other exclusion criteria related to the patient's disease status and level of physical and cognitive function. The cognitive training study included only patients with a KPS between 80 and 100%, without aphasia and without evidence of tumour recurrence after having received standard oncological treatment.<sup>22</sup> A high performance status was also a criterion for inclusion in the massage study and the SN follow-up.<sup>27–28–30</sup> Patients with communication or cognitive problems were excluded from the NTF study,<sup>26</sup> and patient/caregiver groups were excluded if expected survival was less than 3 months.<sup>29–30</sup> Three studies reduced risk of bias by applying a RCT design<sup>29</sup> or by including a control group with a one-to-one match.<sup>24–25</sup> One of these studies excluded subtypes of BT in order to maintain a homogeneous group of participants.<sup>25</sup> The remaining six studies were conducted without a control group.<sup>22–23–26–28–30</sup> Spetz *et al*<sup>27–28</sup> concluded that a SN *function* relies on the person's experience and personal qualities and therefore the function could not be regarded as separate from the *person* performing it, and Sardell *et al*<sup>26</sup> concluded that the study participants only reflected a selected group of patients.

Attrition rate was low for all studies ranging from 0% to 12.5%, except Boele *et al*'s psycho-education intervention reported 51.6% and 32% attrition in the intervention and control groups, respectively. Boele *et al* concluded that a six-session intervention with a psychologist and/or four questionnaires at five test points may have been too demanding for the caregivers. Attrition was not reported in the home training study.<sup>23</sup> An adverse event was reported in the massage

study as one patient developed shingles during the intervention and dropped out.<sup>30</sup> An investigation revealed that this event was unrelated to the massage intervention. Sardell *et al*<sup>26</sup> argued that it was reasonable to assume that HGG patients can be followed up by telephone without adversely affecting overall care or survival. Spetz *et al*<sup>27</sup> elaborated on the risk of the SN being too personally engaged in the relationship with the families that may negatively affect the relationship within the family. Moreover, the authors report that two caregivers abstained from participating in the last interview, which could reflect caregiver dissatisfaction with the intervention.<sup>28</sup> There were no contraindications to physical training reported.<sup>25</sup> The great diversity of age, gender and educational background in the small sample size enabled patients to avoid stressful comparisons and competition in the cognitive training programme, and no adverse events were reported from the cognitive training study.<sup>22</sup> Three studies did not report adverse events.<sup>23–24–29</sup>

## DISCUSSION

This systematic review identified the relatively few controlled trials of unidisciplinary or multidisciplinary non-pharmacological rehabilitative and/or supportive care interventions in patients with HGG and/or their caregivers. The eight studies were found to be diverse in purpose, study design and of low methodological quality. Methodological limitations of the reviewed studies include small sample sizes, biases in study design and few subgroup analyses. However, there is initial evidence that cognitive and physical training and psychosocial support benefit HGG patients and/or their caregivers.

### Rehabilitative interventions

#### Cognitive training

Hassler *et al*<sup>22</sup> found that a neurocognitive group training programme improved memory skills. Late-stage patients with a KPS of 80–100 were enrolled, and it is yet to be explored whether such a programme can be beneficial to HGG patients with a KPS  $\leq 80$ . No control group or follow-up measurements after 12 weeks were conducted, leaving the longitudinal effects of cognitive training unknown among HGG patients.

Based on previous research, there is preliminary evidence to suggest that cognitive training among patients with low-grade gliomas (LGG) improves cognitive status.<sup>31</sup> This RCT study among patients with LGG (N=140) provides initial evidence of a salutary effect on short-term cognitive complaints and on longer-term cognitive performance and mental fatigue.<sup>31</sup> Further, an internet-based cognitive rehabilitation programme in patients with BT proved to be a convenient and flexible alternative to in-clinic programmes.<sup>32</sup>

## Physical training

This systematic review found that early and individually tailored physical training (within weeks after surgery) resulted in significant functional gain for HGG patients. However, functional and neurological gains in the weeks after surgery may be explained by the neurosurgical intervention alone. Geler-Kulcu *et al*'s<sup>24</sup> findings were inconclusive regarding the effect of physical training and occupational therapy in HGG patients as only nine patients with HGG were included.

No physical training studies reported adverse events as a direct result of testing or training. In two studies, functional outcomes were not significantly different in patients with benign tumours versus malignant tumours or primary versus metastatic BTs or BT patients compared with patients with stroke.<sup>23 24</sup> However, in a third study, meningioma patients achieved a greater functional gain (independence in activities of daily living and mobility) compared with patients with HGG or stroke.<sup>25</sup> Evidence for the benefit of physical training among HGG patients is emerging. However, this evidence justifies physical training before the adjuvant therapy as two out of three studies included patients before radiotherapy and chemotherapy was administered.<sup>24 25</sup> Three retrospective studies conclude that patients with BT achieve improved functional outcomes from inpatient rehabilitation.<sup>33–35</sup> Consistent with this, a comparative study of inpatient rehabilitation among patients with BT and patients with traumatic brain injury patients showed functional gains for both groups.<sup>36</sup> However, the absolute functional outcomes were smaller in the BT group.

## Supportive care interventions

## Psychosocial support

In this systematic review, patient and caregivers' need for psychosocial support was met by applying different psychosocial approaches. Psychosocial support provided by a SN is of value for both patients and their families,<sup>27 28</sup> and telephone follow-up is a useful strategy.<sup>26</sup> Repeated counselling with a psychologist improved QOL issues and mastery of caregivers.<sup>29</sup> However, high attrition indicates that the programme was too burdensome for the participants. Keir showed that massage therapy reduces the feeling of stress.<sup>30</sup> It is not possible to estimate the effect of the massage programme as there was no control group. It is imperative that researchers are familiar with the needs, preferences and interests of these patients and their caregivers in order to design interventions that would result in high adherence and low attrition. Two previous reviews explored the needs of patients with HGG and their family/caregivers,<sup>37 38</sup> and presently more evidence is being established.<sup>39</sup>

## Methodological discussion

The *strengths* of the reviewed studies were that patients with HGG and/or their caregivers generally

accept participation in interventions at various times during the disease and treatment trajectory. Interventions that aim to support the patient and/or caregiver, improve cognitive or functional status and reduce stress have high adherence. However, as participation can be experienced as burdensome and result in high attrition, careful ethical and methodological considerations are required. It has been reported that attrition rates are usually high for supportive care interventions in patients with HGG.<sup>40</sup> Reasons for attrition were death, medical decline of the patient or high caregiver burden.<sup>41</sup> High attrition was noted in the RCT that required the participants to respond to four different questionnaires every second month, five times in total.<sup>29</sup> However, the reason for high attrition in this RCT was not confirmed by the participants. Two studies were matched case-control studies<sup>24 25</sup> strengthening the evidence of multidisciplinary interventions (combined physical training with either occupational or speech therapy) on functional outcomes. Further, strengths of the reviewed studies include the use of validated measurement outcomes, except in the qualitative study.<sup>27 28</sup> However, the use of different questionnaires and measurements makes it difficult to compare results.

The *limitations* of the reviewed studies are that they are heterogeneous in many respects, which makes them non-comparable. Sample sizes are small with some studies including as few as 11 participants.<sup>22</sup> This questions the validity and reliability of the statistical analyses in the studies. Further study limitations include lack of control groups and longitudinal follow-up periods. Selection bias arises from the exclusion of patients with low-performance status<sup>22 27 28 30</sup> or cognitive deficit and communication problems.<sup>26</sup>

The scoring system applied for methodological quality needs to be further tested for reliability.<sup>21</sup> However, a pilot study found it to be a promising appraisal tool<sup>42</sup> and it is presently used internationally.<sup>43</sup> Moreover, it provides transparency in how the methodological quality was scored and readers can review the decisions. Finally, an additional descriptive methodological quality appraisal of the included studies supplemented this tool.

## CLINICAL AND RESEARCH IMPLICATIONS

Patients with HGG differ from other cancer diagnoses because of progressive neurological deterioration resulting in changing and complex symptoms along the disease and treatment trajectory. A multidisciplinary approach has the potential to meet the variety of needs emerged from this life-threatening situation. However, cognitive deficits, changes in personality, lack of insight and loss of empathy may impair HGG patients' ability to participate actively in rehabilitative and supportive care programmes. Therefore, it is important that the interventions are tailored to the

special needs of this group. New approaches, such as internet-based approaches, might have the potential to meet the needs of patients with HGG and their caregivers.<sup>44</sup> It is well known that decreased QOL is associated with depression among patients with HGG.<sup>45</sup> However, psychosocial support for patients with HGG has not been the subject of much research,<sup>46</sup> which is why future research should aim at maintaining QOL. Studies that enrol patients during and after oncological treatments are important as the adjuvant treatment has a negative impact on functional capacity among patients with HGG<sup>47</sup> and reduced cardio-respiratory fitness postoperatively.<sup>48</sup> Further, treatment with glucocorticoids causes muscle atrophy.<sup>49</sup> From non-cancer clinical populations, it has shown that resistance and endurance physical training can reverse muscle atrophy.<sup>50</sup> It is therefore interesting to explore whether a similar effect can be achieved for patients with HGG.

## CONCLUSION

As evidence is beginning to emerge, there is a need for well-designed longitudinal and randomised controlled trials of non-pharmacological interventions in patients with HGG and their caregivers. Despite the methodological limitations of the included studies, cognitive training, individual tailored physical training programmes, psychosocial support/education and massage therapy have positive outcomes for both patients and their caregivers, are feasible and so far, without risk for this group. Research is needed to sustain this beginning evidence. To provide stronger evidence, clinical controlled trials with longitudinal designs focusing on cognitive and physical training as well as supportive care are needed.

**Acknowledgements** The authors would like to thank the information specialist Anders Larsen from The University Hospitals Centre for Health Research (UCSF), Copenhagen, Denmark, for his contribution to the literature search.

**Funding** The study was supported by grants from The Centre for Integrated Rehabilitation of Cancer Patients (CIRE), a centre established and supported by The Danish Cancer Society and The Novo Nordisk Foundation. Furthermore, by The Neuro Centre at Copenhagen University Hospital; Rigshospitalet in Denmark, The Novo Nordisk Foundation for Clinical Nursing Research, The Capital Regional Research Foundation in Denmark, Torben and Alice Frimodts Foundation, Vera and Flemming Westerbergs Foundation, Hetland Olsen's Foundation and The Research Foundation of Rigshospitalet, Copenhagen University Hospital.

**Competing interests** None.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Open Access** This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 3.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/3.0/>

## REFERENCES

- 1 Starkweather AR, Sherwood P, Lyon DE, *et al.* A biobehavioral perspective on depressive symptoms in patients with cerebral astrocytoma. *J Neurosci Nurs* 2011;43:17–28.
- 2 Stupp R, Hegi ME, Mason WP, *et al.* Effects of radiotherapy with concomitant and adjuvant temozolomide versus radiotherapy alone on survival in glioblastoma in a randomised phase III study: 5-year analysis of the EORTC-NCIC trial. *Lancet Oncol* 2009;10:459–66.
- 3 Armstrong TS, Cohen MZ, Eriksen LR, *et al.* Symptom clusters in oncology patients and implications for symptom research in people with primary brain tumors. *J Nurs Scholarsh* 2004;36:197–206.
- 4 Kosteljanetz M. Brain tumors—mainly glial tumors in adults. *Ugeskr Laeger* 2002;164:3017–20.
- 5 Molassiotis A, Wilson B, Brunton L, *et al.* Symptom experience in patients with primary brain tumours: a longitudinal exploratory study. *Eur J Oncol Nurs* 2010;14:410–16.
- 6 Lovely MP, Miaskowski C, Dodd M. Relationship between fatigue and quality of life in patients with glioblastoma multiforme. *Oncol Nurs Forum* 1999;26:921–5.
- 7 Kim YH, Park CK, Kim TM, *et al.* Seizures during the management of high-grade gliomas: clinical relevance to disease progression. *J Neurooncol* 2013;113:101–9.
- 8 Klein M, Postma TJ, Taphoorn MJ, *et al.* The prognostic value of cognitive functioning in the survival of patients with high-grade glioma. *Neurology* 2003;61:1796–8.
- 9 Boele FW, Klein M, Reijneveld JC, *et al.* Symptom management and quality of life in glioma patients. *CNS Oncol* 2014;3:37–47.
- 10 Davies E, Higginson IJ. Communication, information and support for adults with malignant cerebral glioma: a systematic literature review. *Support Care Cancer* 2003;11:9–21.
- 11 Diaz JL, Barreto P, Gallego JM, *et al.* Proper information during the surgical decision-making process lowers the anxiety of patients with high-grade gliomas. *Acta Neurochir (Wien)* 2009;151:357–62.
- 12 Lobb EA, Halkett GK, Nowak AK. Patient and caregiver perceptions of communication of prognosis in high grade glioma. *J Neurooncol* 2011;104:315–22.
- 13 Rooney AG, McNamara S, Mackinnon M, *et al.* The frequency, longitudinal course, clinical associations, and causes of emotional distress during primary treatment of cerebral glioma. *Neuro Oncol* 2013;15:635–43.
- 14 Keir ST. Levels of stress and intervention preferences of caregivers of brain tumor patients. *Cancer Nurs* 2007;30:E33–9.
- 15 Catt S, Chalmers A, Fallowfield L. Psychosocial and supportive-care needs in high-grade glioma. *Lancet Oncol* 2008;9:884–91.
- 16 Khan F, Amatya B, Ng L, *et al.* Multidisciplinary rehabilitation after primary brain tumour treatment. *Cochrane Database Syst Rev* 2013;1:CD009509.
- 17 Halkett GK, Lobb EA, Oldham L, *et al.* The information and support needs of patients diagnosed with high grade glioma. *Patient Educ Couns* 2010;79:112–19.
- 18 Parvataneni R, Polley MY, Freeman T, *et al.* Identifying the needs of brain tumor patients and their caregivers. *J Neurooncol* 2011;104:737–44.
- 19 Liberati A, Altman DG, Tetzlaff J, *et al.* The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: Explanation and elaboration. *BMJ* 2009;339:b2700.



- 20 Skorkjær Binderkrantz A, Bøgh Andersen L. *Guide to NVivo 9*. 1st edn. Hans Reitzels, 2011.
- 21 Pluye P, Gagnon MP, Griffiths F, *et al*. A scoring system for appraising mixed methods research and concomitantly appraising qualitative, quantitative and mixed methods primary studies in mixed studies reviews. *Int J Nurs Stud* 2009;46:529–46.
- 22 Hassler MR, Elandt K, Preusser M, *et al*. Neurocognitive training in patients with high-grade glioma: A pilot study. *J Neurooncol* 2010;97:109–15.
- 23 Pace A, Parisi C, Di Lelio M, *et al*. Home rehabilitation for brain tumor patients. *J Exp Clin Cancer Res* 2007;26:297–300.
- 24 Geler-Kulcu D, Gulsen G, Buyukbaba E, *et al*. Functional recovery of patients with brain tumor or acute stroke after rehabilitation: a comparative study. *J Clin Neurosci* 2009;16:74–8.
- 25 Bartolo M, Zucchella C, Pace A, *et al*. Early rehabilitation after surgery improves functional outcome in inpatients with brain tumours. *J Neurooncol* 2012;107:537–44.
- 26 Sardell S, Sharpe G, Ashley S, *et al*. Evaluation of a nurse-led telephone clinic in the follow-up of patients with malignant glioma. *Clin Oncol (R Coll Radiol)* 2000;12:36–41.
- 27 Spetz A, Henriksson R, Bergenheim AT, *et al*. A specialist nurse-function in neurooncology: a qualitative study of possibilities, limitations, and pitfalls. *Palliat Support Care* 2005;3:121–30.
- 28 Spetz A, Henriksson R, Salander P. A specialist nurse as a resource for family members to patients with brain tumors: An action research study. *Cancer Nurs* 2008;31:E18–26.
- 29 Boele FW, Hoebe W, Hilverda K, *et al*. Enhancing quality of life and mastery of informal caregivers of high-grade glioma patients: a randomized controlled trial. *J Neurooncol* 2013;111:303–11.
- 30 Keir ST. Effect of massage therapy on stress levels and quality of life in brain tumor patients-observations from a pilot study. *Support Care Cancer* 2011;19:711–5.
- 31 Gehring K, Sitskoorn MM, Gundy CM, *et al*. Cognitive rehabilitation in patients with gliomas: a randomized, controlled trial. *J Clin Oncol* 2009;27:3712–22.
- 32 Gehring K, Aaronson NK, Taphoorn MJ, *et al*. Interventions for cognitive deficits in patients with a brain tumor: An update. *Expert Rev Anticancer Ther* 2010;10:1779–95.
- 33 Greenberg E, Tregler I, Ring H. Rehabilitation outcomes in patients with brain tumors and acute stroke: Comparative study of inpatient rehabilitation. *Am J Phys Med Rehabil* 2006;85:568–73.
- 34 Fu JB, Parsons HA, Shin KY, *et al*. Comparison of functional outcomes in low- and high-grade astrocytoma rehabilitation inpatients. *Am J Phys Med Rehabil* 2010;89:205–12.
- 35 Marciniak CM, Sliwa JA, Heinemann AW, *et al*. Functional outcomes of persons with brain tumors after inpatient rehabilitation. *Arch Phys Med Rehabil* 2001;82:457–63.
- 36 O'Dell MW, Barr K, Spanier D, *et al*. Functional outcome of inpatient rehabilitation in persons with brain tumors. *Arch Phys Med Rehabil* 1998;79:1530–4.
- 37 Sterckx W, Coolbrandt A, Dierckx de Casterle B, *et al*. The impact of a high-grade glioma on everyday life: A systematic review from the patient's and caregiver's perspective. *Eur J Oncol Nurs* 2013;17:107–17.
- 38 Moore G, Collins A, Brand C, *et al*. Palliative and supportive care needs of patients with high-grade glioma and their carers: A systematic review of qualitative literature. *Patient Educ Couns* 2013;91:141–53.
- 39 Piil K, Jarden M, Jakobsen J, *et al*. A longitudinal, qualitative and quantitative exploration of daily life and need for rehabilitation among patients with high-grade gliomas and their caregivers. *BMJ Open* 2013;3(7). pii: e003183.
- 40 Ford E, Catt S, Chalmers A, *et al*. Systematic review of supportive care needs in patients with primary malignant brain tumors. *Neuro Oncol* 2012;14:392–404.
- 41 Meyers FJ, Carducci M, Loscalzo MJ, *et al*. Effects of a problem-solving intervention (COPE) on quality of life for patients with advanced cancer on clinical trials and their caregivers: Simultaneous care educational intervention (SCEI): Linking palliation and clinical trials. *J Palliat Med* 2011;14:465–73.
- 42 Pace R, Pluye P, Bartlett G, *et al*. Testing the reliability and efficiency of the pilot mixed methods appraisal tool (MMAT) for systematic mixed studies review. *Int J Nurs Stud* 2012;49:47–53.
- 43 Polit D, Beck C, eds. *Nursing research. Generating and assessing evidence for nursing practice*. 9th edn. Wolters Kluwer. Lippincott Williams & Wilkins, 2012.
- 44 Jung Oh H, Ozkaya E, LaRose R. How does online social networking enhance life satisfaction? The relationships among online supportive interaction, affect, perceived social support, sense of community, and life satisfaction. *Comput Hum Behav* 2014;30:69–78.
- 45 Cheng JX, Zhang X, Liu BL. Health-related quality of life in patients with high-grade glioma. *Neuro Oncol* 2009; 11:41–50.
- 46 Pace A, Metro G, Fabi A. Supportive care in neurooncology. *Curr Opin Oncol* 2010;22:621–6.
- 47 Jones LW, Guill B, Keir ST, *et al*. Exercise interest and preferences among patients diagnosed with primary brain cancer. *Support Care Cancer* 2007;15:47–55.
- 48 Jones LW, Friedman AH, West MJ, *et al*. Quantitative assessment of cardiorespiratory fitness, skeletal muscle function, and body composition in adults with primary malignant glioma. *Cancer* 2010;116:695–704.
- 49 Hempen C, Weiss E, Hess CF. Dexamethasone treatment in patients with brain metastases and primary brain tumors: Do the benefits outweigh the side-effects? *Support Care Cancer* 2002;10:322–8.
- 50 Horber FF, Hoopeler H, Scheidegger JR, *et al*. Impact of physical training on the ultrastructure of mid thigh muscle in normal subjects and in patients treated with glucocorticoids. *J Clin Invest* 1987;79:1181–90.



Supplementary Table 1 Description of the studies

Reference and country	Population, sample size and attrition	Median age (range)	Study design, focus and time of enrollment	Intervention, mode and setting	Duration, frequency, length and adherence	Time for measurements and main results
Hassler et al., 2010 <sup>22</sup>  Austria	HGG (n=11)  Attrition 0%	50 years (range 23-73)	Clinical controlled pilot trial, single arm, no control group  Investigates the outcome on cognitive function  Enrollment 10-42 months (median 15) after diagnosis	Neuro-cognitive group training  Group + supervised  Setting: outpatient clinic	1½ hour/1 day per week for 10 weeks  100% adherence and satisfaction	Pre- and post measurements  Verbal Memory Total Learning (HVLTL total learning) ↑ ( $P=0.04$ )  Verbal Memory Delayed Recall (HVLTL recall) → ( $P=0.11$ )  Psychomotor speed (TMT-A) → ( $P=0.22$ )  Sustained attention (TMT-B) → ( $P=0.17$ )  Verbal fluency (COWAT) → ( $P=0.29$ )
Pace et.	BT	60	Clinical	Physical	1 hour/ 3	Pre- and post

Supplementary Table 1 Description of the studies

al. 2007 <sup>23</sup>	(121); 62 with HGG, 27 with AA, 12 with Oligode n- droglio ma and mixed BT, 20 with other BT.  Attrition not reported	years (range 14-79)	controlled pilot trial, single arm, no control group  Explores the functional outcome and impact on QOL  Enrollment: 82 after diagnosis, 39 after recurrence	training  Individual + supervised  Setting: home	days per week for 12 weeks  Adherence not reported	measurements  BI: An increased tendency in 39%, a stable tendency in 16%, and decreased tendency in 44% (no <i>P</i> value)  A tendency for increased KPS in 24% (no <i>P</i> value)  Among a selected group of clinical improved patients: Median BI↑ ( <i>P</i> =0.001) and Median KPS ↑ ( <i>P</i> = 0.001)  No differences in functional gain between the histological subgroup (no <i>P</i> value)
------------------------	---	---------------------------	--	--	--	---

Supplementary Table 1 Description of the studies

						<p>A functional gain was observed in 26% of the patients with HGG (not specified)</p> <p>72% had an improvement in at least one domain scores compared with their baseline QOL score (n=54)</p>
<p>Geler- Kulcu et al. 2008 <sup>24</sup>  Turkey</p>	<p>BT (n=21); 9 with HGG Stroke (n=21)     Attrition 0%</p>	<p>BT group: 52.5 (range 36.5- 68.5)  Stroke group: 56.7 (range 45.7- 67.7)</p>	<p>Clinical controlled and comparative trial, single arm, two groups matched according to lesion side  Examines the functional</p>	<p>Physical training + Occupation al therapy  Individual + supervised  Setting: hospital</p>	<p>1 hour/5 days per week for 5.7- 18.7 weeks  Mean number of sessions: BT group 11.5± 5.8 Stroke group 12.3±6.4</p>	<p>Pre- and post measurements  Both the BT group and the stroke group improved significantly:  MAS ↑ (<math>P=0.025/ P =</math> 0.002) between groups (<math>P=0.718</math>) PASS ↑ (<math>P=0.002/</math> <math>P=0.002</math>) between groups (<math>P=0.817</math>) BBS ↑ (<math>P=0.011/</math></p>

Supplementary Table 1 Description of the studies

			outcomes		100% adherence	<p><math>P=0.109</math>) between groups (<math>P=0.109</math>)</p> <p>FIM mobility ↑ (<math>P=0.043/ P=0.007</math>) between groups (<math>P=0.660</math>)</p> <p>There was no statistically significant difference between the two groups in any of the four outcome measures.</p>
Bartolo et al. 2012 <sup>25</sup>  Italy	BT (n=75); 43 with HGG, 32 with meningioma Stroke (n=75)  Attrition	BT group: 63.5 (range 50.3-76.7)  HGG group: 62 (range	A case-controlled and comparative trial, single arm, two groups matched one-to-one, for age, sex, and side of	Individual + supervised  Physical training + speech therapy  Setting: Hospital	1 hour/ 6 days a week for 4 weeks  100% adherence	<p>Pre- and post measurements</p> <p>FIM, Sitting Balance, Standing balance, Hauser Index and MGHFAC ↑ for both groups (<math>P&lt;0.001</math>)</p> <p>Patients with meningioma achieved</p>



Supplementary Table 1 Description of the studies

	0%	48.1-75.9)	lesion			better results in regards to independence in activities of daily living ( $P=0.02$ ) and mobility ( $P=0.04$ ) compared with HGG or stroke.
		Stroke group: 65.7 (range 55.1-76.3)	Tests the functional outcomes  Enrollment after surgery			
Sardell et al., 2000 <sup>26</sup>  UK	BT (N=45); 43 with HGG, 1 with PNET, 1 with OA  Attrition: unclear	50 years (range 23-69)	A quantitative pilot evaluation study  Measures the numbers of unscheduled outpatient visits, BI, and patient satisfaction	Individual + supervised  Psycho-social and supportive intervention with telephone follow-up in a nurse-led telephone follow-up clinic.	1 telephone consultation/month for 3 months + face-to-face consultation in the clinic on the fourth month. Patients	Unscheduled clinic visits and BI: pre and post measurements  Satisfaction questionnaire completed at the fourth month or later (n=22)  Unscheduled clinic visits= 41. Thirty-one were due to disease progression and were usually initiated by the nurse after a NTF call.

Supplementary Table 1 Description of the studies

			(n=22)  Enrollment 16 months after diagnosis	Setting:  Combined home and outpatient	could  choose to continue with NTF hereafter  Adherence : 80% after 4 month, 45% at 8 months, 34% at 1 year.	The other 10 visits were due to concern regarding symptoms.  The number of unscheduled clinic visits would not have been avoided with CCF carried out at the same time intervals.  BI at baseline: 43 patients reported normal or near normal function. On subsequent follow-up (n=11) and time not specified – 5 patients had an improvement, six had a reduction, one patient was not assessed (no <i>P</i> -value)  Patient satisfaction: 81% strongly agree and 19% agree that
--	--	--	---	--	---	---

Supplementary Table 1 Description of the studies

						the follow up allows the patient to get on with their life and make better use of their time. Median patient satisfaction score of 9 on a visual analogue scale of 0-10 (range 3.6-10)
Spetz et al., 2005 <sup>27</sup>  Sweden	HGG (n=16)  Caregivers (n=16);  10 spouses       Attrition: 0%	Patients: 50.8 (range from 21-71)  Caregivers: not reported	A qualitative descriptive intervention study  Explores the patients' and their families' perspective on the function, relationship and their use of a SN	Individual + supervised  Psycho-social and supportive intervention with a SN function  Setting: Combined home and outpatient	Ongoing telephone contact with a SN during the entire course.       100% adherence	Field notes and conversations were written during the entire course of the disease. Interviews with patients and their families every third month. The spouses were interviewed after the patient's death  Four relationships between the family and the SN were identified:  1) a resource for the

Supplementary Table 1 Description of the studies

			function  Enrollment at time of diagnosis			family 2) a parallel resource within the family 3) a resource for the caregiver 4) a resource for the patient
Spetz et al., 2008 <sup>28</sup>  Sweden	HGG patients (n=16)  Caregiv ers (n=16)  Attrition: 0%  Attrition from the last interview: 12.5%	Same study as describ ed in Spetz, 2005 <sup>26</sup>	A qualitative descriptive intervention study  Explores how the caregivers utilized the SN function and the relationship between the SN and the caregivers  Enrollment at time of diagnosis	Same study as described in Spetz, 2005 <sup>26</sup>	Same study as described in Spetz, 2005 <sup>26</sup>	Field notes and conversations were written during the entire course of the disease. Interviews with patients and their families every third month. The spouses were interviewed after the patient's death  Mean number of total contacts= 32 (range 1- 115)  Three main categories of subjects discussed with the SN were: 1) Subjects related to the



Supplementary Table 1 Description of the studies

						<p>sick partner's medical health and treatment</p> <p>2) Subjects related to needs and desires concerning one-self 3) Subjects related to the relationship with the SN</p> <p>Close relationships between the SN and the caregivers were identified as reducing the caregivers' feeling of being left alone with the responsibility, and assisted adequate decisions</p>
Boele et al., 2011 <sup>29</sup>  The	IG: caregivers (n=31)  CG: caregiver	IG: 50.77 (range 39.3-62.2)	A cross-sectional randomized controlled intervention trial	Individual+ supervised  Psycho-social and	IG: 1 hour/1 day per week for 6 weeks  CG: Usual	<p>Four questionnaires at baseline and every 2 months, five times in total</p> <p>Patients increased</p>

Supplementary Table 1 Description of the studies

Netherlands	rs (n=25)  Attrition: IG: 51.6% CG: 32%	CG: 50.56 (range 40.2 - 60.92)	Investigates the outcome on QOL and feeling of mastery of caregivers  Enrollment ≥12 months from diagnosis (n=24) ≤12 months from diagnosis (n=32)	educational intervention  Setting: Hospital	care e.g. interaction s with specialists  Adherence : IG:48.4% CG:68%	uncertainty concerning the future and increments in communication deficits were negatively associated with mental functioning of the caregiver ( $P=0.004$ and $P=0.043$ , respectively)  Caregiver mastery was positively related to patient's visual disorders ( $P=0.031$ ) + communication deficits ( $P=0.005$ )  Caregiver mastery delta scores (assessment at test time 5 minus baseline assessment) differed significantly between
-------------	--	--	--	---	---	---

Supplementary Table 1 Description of the studies

						the two groups ( $P=0.008$ ) showing that the IG's feelings of mastery increased ( $P=0.021$ )
Keir, 2011 <sup>30</sup>  USA	HGG (n=26) with KPS of $\geq 70$ .  Attrition: one patient dropped out	Not reported	A prospective, single-arm intervention pilot study, no control group  Assesses the patient reported psychologic al outcomes, level of stress and QOL  Enrollment	Individual+ supervised  Massage therapy  Setting: A outpatient clinic related to the Hospital	45 minutes/2 days per week for 4 weeks  Adherence :96.2%	Measurements at baseline, after week 1,2,3, and 4 and 1 week after last massage session (week 5) Levels of stress ↓ between week 2 and 3 ( $P=0.010$ ) A tendency for reduction in stress continued throughout week 4 ( $P=0.063$ ) FACT-Br at week 4: Emotional well-being ↑ ( $P\leq 0.001$ ) Additional BT concerns ↑ ( $P\leq 0.013$ ) Social/family well-

Supplementary Table 1 Description of the studies

			after diagnosis			being ↑ ( $P \leq 0.024$ )  A trend for improvement in physical well-being ( $P \leq 0.078$ )  At the end of week 4, PSS-10 scores of all participants were below the stress threshold
--	--	--	--------------------	--	--	---

Abbreviations: HGG, Patients diagnosed with a high-grade glioma; KPS, Karnofsky Performance Score; AA, Anaplastic Astrocytoma; BT, Brain tumor; TMT-B, Trail Making Test B; TMT-A, Trail Making Test A; COWAT, Controlled Oral Word Association Test; BI, Barthel Index used to measure Activities of Daily Living; MAS, Motor Assessment Scale; PASS, Postural Assessment Scale for Stroke; BBS, Berg Balance Scale; FIM, Functional Impairment Measurement; MGHFAC, Massachusetts General Hospital Functional Ambulation Classification; PNET, Primitive neuroectodermal tumor; OA, Oligoastrocytoma; SN, Specialist Nurse; NTF, Nurse-led Telephone Follow-up; CCF, Conventional Clinic Follow-up; PSS-10, Perceived Stress Scale; FACT-Br, Functional Assessment of Cancer Therapy-Brain; IG, Intervention group; CG, Control group; QOL, Quality of life; →, No change; ↑, Significant increased change; ↓, Significant decreased change.



Supplementary Table 2 The methodological quality

Type of study	Methodological quality criteria	Hassler et al. 2010 <sup>22</sup>	Pace et al. 2007 <sup>23</sup>	Geler-Kulcu et al. 2008 <sup>24</sup>	Bartolo et al. 2012 <sup>25</sup>	Sardell et al. 2000 <sup>26</sup>	Spetz et al. 2005 <sup>27</sup>	Spetz et al. 2008 <sup>28</sup>	Boele et al. 2012 <sup>29</sup>	Keir 2011 <sup>30</sup>
Qualitative	Qualitative objective or question	-	-	-	-	-	1	1	-	-
	Appropriate qualitative approach or design or method	-	-	-	-	-	1	1	-	-
	Description of the	-	-	-	-	-	1	1	-	-

Supplementary Table 2 The methodological quality

	context									
	Descri ption of particip ants and justific ation of sampli ng	-	-	-	-	-	1	1	-	-
	Descri ption of qualitat ive data collecti on and analysi s	-	-	-	-	-	1	1	-	-
	Discus sion of	-	-	-	-	-	1	1	-	-

Supplementary Table 2 The methodological quality

	researchers' reflexivity									
Quantitative Experimental	Appropriate sequence generation and/or randomization	0	0	0	0	0	-	-	1	0
	Allocation concealment and/or blinding	0	0	0	0	0	-	-	0	0
	Complete outcome	1	1	1	1	1	-	-	0	1





Supplementary Table 2 The methodological quality

	on- analysi s techniq ues or proced ures									
	Integra tion of qualitat ive and quantit ative data or results	-	-	-	-	-	-	-	-	-
Sco re in per cent		33,3%	33,3 %	33,3%	33,3 %	33,3 %	100%	100%	33,3 %	33,3 %