

HEART FAILURE CARE OR PALLIATIVE CARE? PATIENTS WANT AND NEED BOTH

¹Mary Brice, ²Amanda Martin, ¹Penny Hansford, ¹Heather Richardson, ^{1,3}Rob George, ³Fliss Murtagh. ¹St Christopher's Hospice, UK; ²Central and Northwest London NHS Foundation Trust, UK; ³King's College London, UK

10.1136/bmjspcare-2016-001204.10

Introduction People with advanced heart failure have significant symptom/disease burden. Optimal models of care have not yet been widely tested and evaluated. We aimed to test a novel combined heart failure palliative care service model.

Method A mixed-method feasibility study, including comparison with an age- and gender-matched cohort receiving specialist palliative care.

Results 27 male and 13 female patients (mean age 84 years, range 60–99) were recruited. Integrated Palliative care Outcome Scale scores, compared with an age- gender- matched specialist palliative care population, demonstrated significantly higher levels of fatigue (mean score 2.6 v 1.9, $p = 0.01$), poor mobility (2.75 v 1.6, $p = 0.00$), drowsiness (2.03 v 0.6, $p = 0.00$), breathlessness (2.2 v 0.9, $p = 0.00$), sore/dry mouth (1.4 v 0.6, $p = 0.004$), depression (1.4 v 0.5, $p = 0.002$) and not feeling at peace (1.6 v 0.9, $p = 0.02$). 67% (95% CI: 49–79%) were severely/overwhelmingly affected by poor mobility and 50% (95% CI: 34–65%) were affected by weakness/fatigue in the previous week. In qualitative interviews (8 patients, 6 carers and 2 joint), patients sought heart failure care critical for optimal symptom control, and whilst realistic about the limitations of therapies, wanted parallel support to prepare for and manage future deterioration.

Conclusion People with advanced heart failure have considerably greater needs than an age-matched palliative care population. They have realistic expectations about interventions available but identify practical support and appropriately skilled healthcare teams as important to their care. Professionals skilled in both heart failure and palliative care interventions in an integrated service model are needed.

REFERENCES

- 1 Hogg KJ, Jenkins SMM. Prognostication or identification of palliative needs in advanced heart failure: where should the focus lie? *BMJ* 2012;**98**(7):523524
- 2 Nordgren L, Soerensen S. Symptoms experienced in the last six months of life in patients with end-stage heart failure. *Eur J Cardiovasc Nurs* 2003;**2**(1):213217
- 3 Pattenden JF, Mason AR, Lewin RJP. Collaborative palliative care for advanced heart failure: outcomes and costs from the 'Better Together' pilot study. *BMJ Support Palliat Care* 2013;**3**(1):69–76
- 4 Selman L, Harding R, Beynon T, *et al.* Modelling services to meet the palliative care needs of chronic heart failure patients and their patients: current practice in the UK. *Palliat Med* 2007;**21**(5):385–390